EP200

FUNCTIONAL FERMENTED PROTEIN

DATASHEET Version: 18-05-2021

EP200 is a complementary feed produced from soybean meal fermented with lactic acid bacteria. EP200 contains fermentation metabolites, including lactic acid, and contains gut flora stabilizing lactic acid bacteria. The fermentation process makes the product highly digestible. EP200 is suitable for young and vulnerable animals and helps maintaining a healthy and functional gut.

NUTRIENTS

Dry Matter	900	g/kg
Crude Protein	505	g/kg
Crude Fat	20	g/kg
Crude Fibre	35	g/kg
Crude Ash	65	g/kg
Starch	30	g/kg
Sugars	30	g/kg

P			
Gross	SID Pigs	SID Poultry	
30.7	27.0	27.3	g/kg
7.2	6.6	6.7	g/kg
14.3	12.6	13.3	g/kg
21.5	19.2	20.2	g/kg
7.0	6.4	6.7	g/kg
23.9	21.7	22.5	g/kg
37.1	35.3	35.9	g/kg
13.9	12.4	13.1	g/kg
40.7	37.1	38.6	g/kg
25.1	22.5	23.0	g/kg
27.3	25.1	25.9	g/kg
18.7	17.0	17.6	g/kg
	30.7 7.2 14.3 21.5 7.0 23.9 37.1 13.9 40.7 25.1 27.3	Gross Pigs 30.7 27.0 7.2 6.6 14.3 12.6 21.5 19.2 7.0 6.4 23.9 21.7 37.1 35.3 13.9 12.4 40.7 37.1 25.1 22.5 27.3 25.1	GrossPigsPoultry30.727.027.37.26.66.714.312.613.321.519.220.27.06.46.723.921.722.537.135.335.913.912.413.140.737.138.625.122.523.027.325.125.9

ENERGY

ME Pigs	14.7 MJ/kg
ME Broilers	10.0 MJ/kg
ME Layers	10.4 MJ/kg

MINERALS

Calcium	2.6	g/kg
Phosphorus	7.2	g/kg
Dig. Phosphorus	4.6	g/kg
Sodium	0.3	g/kg
Potassium	21.7	g/kg
Chloride	0.4	g/kg

ANTINUTRITIONAL SUBSTANCES

TIA	< 0.5 mg/g
β-conglycinine	< 0.5 _{mg/g}
Lectins	< 0.05 mg/g
Raffinose and Stachyose	< 0.5 %

LACTIC ACID AND BACTERIA

Lactic acid	60 g/kg
Lactic acid bacteria *	>10 ⁶ cfu/g

* Enterococcus faecium (NCIMB 10415)

EP200 is a fermented complementary feed composed out of plant raw materials. The figures are based on analyses representing an average. Some variance in physico-chemical composition may occur.



Better gut

animals

health for your



Appearance:	Yellowish light brown meal with a pleasant flavor of fermentation
Package size:	Big bags (1000kg) and bags (25kg)
Storage & Shelf life:	12 months from production date when stored at 20°C or below. Shelf life will be shorter if stored at a higher temperature.

Instructions for use*:

	Period	EP200
PIGS		
Piglets	Before and after weaning	8-15%
	12-20 kg	1-8%
POULTRY		
Broilers	Until day 14	5%
	Until day 24	3%
	After day 24	2%
AQUACULTURE		
Salmons		15-25%

* Percent of dry feed.

The product contains 6% lactic acid. Remember to optimize and reduce added acidifiers.