


# EP200 non-GMO

## FUNCTIONAL FERMENTED PROTEIN

DATASHEET

Version: 18-05-2021

EP200 non-GMO is a complementary feed produced from soybean meal fermented with lactic acid bacteria. EP200 non-GMO contains fermentation metabolites, including lactic acid, and contains gut flora stabilizing lactic acid bacteria. The fermentation process makes the product highly digestible. EP200 non-GMO is suitable for young and vulnerable animals and helps maintaining a healthy and functional gut.



**Better gut  
health for your  
animals**

### NUTRIENTS

Dry Matter	900 g/kg
Crude Protein	505 g/kg
Crude Fat	20 g/kg
Crude Fibre	35 g/kg
Crude Ash	65 g/kg
Starch	30 g/kg
Sugars	30 g/kg

### ENERGY

ME Pigs	14.7 MJ/kg
ME Broilers	10.0 MJ/kg
ME Layers	10.4 MJ/kg

### MINERALS

Calcium	2.6 g/kg
Phosphorus	7.2 g/kg
Dig. Phosphorus	4.6 g/kg
Sodium	0.3 g/kg
Potassium	21.7 g/kg
Chloride	0.4 g/kg

### LACTIC ACID AND BACTERIA

Lactic acid	60 g/kg
Lactic acid bacteria *	> 10 <sup>6</sup> cfu/g

\* Enterococcus faecium (NCIMB 10415)

EP200 non-GMO is a fermented complementary feed composed out of plant raw materials. The figures are based on analyses representing an average. Some variance in physico-chemical composition may occur.

AMINO ACIDS	Gross	SID Pigs	SID Poultry	
Lys	30.7	27.0	27.3	g/kg
Met	7.2	6.6	6.7	g/kg
M+C	14.3	12.6	13.3	g/kg
Thre	21.5	19.2	20.2	g/kg
Tryp	7.0	6.4	6.7	g/kg
Ile	23.9	21.7	22.5	g/kg
Arg	37.1	35.3	35.9	g/kg
His	13.9	12.4	13.1	g/kg
Leu	40.7	37.1	38.6	g/kg
Val	25.1	22.5	23.0	g/kg
Phen	27.3	25.1	25.9	g/kg
Tyr	18.7	17.0	17.6	g/kg

### ANTINUTRITIONAL SUBSTANCES

TIA	< 0.5 mg/g
β-conglycinine	< 0.5 mg/g
Lectins	< 0.05 mg/g
Raffinose and Stachyose	< 0.5 %

**Appearance:**

Yellowish light brown meal with a pleasant flavor of fermentation

**Package size:**

Big bags (1000kg) and bags (25kg)

**Storage & Shelf life:**

12 months from production date when stored at 20°C or below.  
Shelf life will be shorter if stored at a higher temperature.

**Instructions for use\*:**

	Period	EP200 non-GMO
<b>PIGS</b>		
<b>Piglets</b>	Before and after weaning	8-15%
	12-20 kg	1-8%
<b>POULTRY</b>		
<b>Broilers</b>	Until day 14	5%
	Until day 24	3%
	After day 24	2%
<b>AQUACULTURE</b>		
<b>Salmons</b>		15-25%

\* Percent of dry feed.

The product contains 6% lactic acid. Remember to optimize and reduce added acidifiers.